

Docket: 13020-10

## AUTOMATIC GENOTYPE DETERMINATION

09648178 071800

Inventors: Stephen E. Lincoln  
Michael R. Knapp

**Certificate of Express Mail Under 37 CFR 1.10**

I hereby declare that this correspondence is being deposited with the United States Postal Service via Express Mail Label No. EL301953995US\_ in an envelope addressed to: Commissioner of Patents and Trademarks, Washington, DC 20231

July 18, 2000  
Date

Evan McHale  
Name

Attorney Docket 1471/108

## AUTOMATIC GENOTYPE DETERMINATION

Cross Reference to Related Applications

*5*  
*Amo A1*  
This application is <sup>a</sup>a continuation of application serial no. 08/362,266, filed December 22, 1994, which is a continuation in part of application serial no. 08/173,173, filed December 23, 1993, which is for an invention entitled "Automatic Genotype Determination," by Stephen E. Lincoln and Michael P. Knapp. This immediate parent application <sup>a2</sup>is a continuation in part of application serial no. 07/775,786, filed October 11, 1991, for an invention entitled "Nucleic Acid Typing by Polymerase Extension of Oligonucleotides using Terminator Mixtures," by P. Goelet, M. Knapp, and S. Anderson, which in turn is a continuation in part of application serial no. 07/664,837, filed March 5, 1991. Immediate parent application serial no. 08/173,173 is also a continuation in part of application serial no. 08/162,397, filed December 6, 1993, for an invention entitled "Method for Immobilization of Nucleic Acid Molecules" by T. Nikiforov and M. Knapp, and of application serial no. 08/155,746, filed November 23, 1993, for an invention entitled "Method for Generating Single-Stranded DNA Molecules" by T. Nikiforov and M. Knapp, and of application serial no. 08/145,145, filed November 3, 1993, for an invention entitled "Single Nucleotide Polymorphisms and their use in Genetic Analysis" by M. Knapp and P. Goelet. All of these related applications are hereby incorporated herein by reference.

5

Technical Field

The present invention relates to the methods and devices for determining the genotype at a locus within genetic material.

10

Summary of the Invention

095415  
095416  
095417  
095418  
095419  
095420  
095421  
095422  
095423  
095424  
095425  
095426  
095427  
095428  
095429  
095430  
095431  
095432  
095433  
095434  
095435  
095436  
095437  
095438  
095439  
095440  
095441  
095442  
095443  
095444  
095445  
095446  
095447  
095448  
095449  
095450  
095451  
095452  
095453  
095454  
095455  
095456  
095457  
095458  
095459  
095460  
095461  
095462  
095463  
095464  
095465  
095466  
095467  
095468  
095469  
095470  
095471  
095472  
095473  
095474  
095475  
095476  
095477  
095478  
095479  
095480  
095481  
095482  
095483  
095484  
095485  
095486  
095487  
095488  
095489  
095490  
095491  
095492  
095493  
095494  
095495  
095496  
095497  
095498  
095499  
095500  
095501  
095502  
095503  
095504  
095505  
095506  
095507  
095508  
095509  
095510  
095511  
095512  
095513  
095514  
095515  
095516  
095517  
095518  
095519  
095520  
095521  
095522  
095523  
095524  
095525  
095526  
095527  
095528  
095529  
095530  
095531  
095532  
095533  
095534  
095535  
095536  
095537  
095538  
095539  
095540  
095541  
095542  
095543  
095544  
095545  
095546  
095547  
095548  
095549  
095550  
095551  
095552  
095553  
095554  
095555  
095556  
095557  
095558  
095559  
095560  
095561  
095562  
095563  
095564  
095565  
095566  
095567  
095568  
095569  
095570  
095571  
095572  
095573  
095574  
095575  
095576  
095577  
095578  
095579  
095580  
095581  
095582  
095583  
095584  
095585  
095586  
095587  
095588  
095589  
095590  
095591  
095592  
095593  
095594  
095595  
095596  
095597  
095598  
095599  
095600  
095601  
095602  
095603  
095604  
095605  
095606  
095607  
095608  
095609  
095610  
095611  
095612  
095613  
095614  
095615  
095616  
095617  
095618  
095619  
095620  
095621  
095622  
095623  
095624  
095625  
095626  
095627  
095628  
095629  
095630  
095631  
095632  
095633  
095634  
095635  
095636  
095637  
095638  
095639  
095640  
095641  
095642  
095643  
095644  
095645  
095646  
095647  
095648  
095649  
095650  
095651  
095652  
095653  
095654  
095655  
095656  
095657  
095658  
095659  
095660  
095661  
095662  
095663  
095664  
095665  
095666  
095667  
095668  
095669  
095670  
095671  
095672  
095673  
095674  
095675  
095676  
095677  
095678  
095679  
095680  
095681  
095682  
095683  
095684  
095685  
095686  
095687  
095688  
095689  
095690  
095691  
095692  
095693  
095694  
095695  
095696  
095697  
095698  
095699  
095700  
095701  
095702  
095703  
095704  
095705  
095706  
095707  
095708  
095709  
095710  
095711  
095712  
095713  
095714  
095715  
095716  
095717  
095718  
095719  
095720  
095721  
095722  
095723  
095724  
095725  
095726  
095727  
095728  
095729  
095730  
095731  
095732  
095733  
095734  
095735  
095736  
095737  
095738  
095739  
095740  
095741  
095742  
095743  
095744  
095745  
095746  
095747  
095748  
095749  
095750  
095751  
095752  
095753  
095754  
095755  
095756  
095757  
095758  
095759  
095760  
095761  
095762  
095763  
095764  
095765  
095766  
095767  
095768  
095769  
095770  
095771  
095772  
095773  
095774  
095775  
095776  
095777  
095778  
095779  
095780  
095781  
095782  
095783  
095784  
095785  
095786  
095787  
095788  
095789  
095790  
095791  
095792  
095793  
095794  
095795  
095796  
095797  
095798  
095799  
095800  
095801  
095802  
095803  
095804  
095805  
095806  
095807  
095808  
095809  
095810  
095811  
095812  
095813  
095814  
095815  
095816  
095817  
095818  
095819  
095820  
095821  
095822  
095823  
095824  
095825  
095826  
095827  
095828  
095829  
095830  
095831  
095832  
095833  
095834  
095835  
095836  
095837  
095838  
095839  
095840  
095841  
095842  
095843  
095844  
095845  
095846  
095847  
095848  
095849  
095850  
095851  
095852  
095853  
095854  
095855  
095856  
095857  
095858  
095859  
095860  
095861  
095862  
095863  
095864  
095865  
095866  
095867  
095868  
095869  
095870  
095871  
095872  
095873  
095874  
095875  
095876  
095877  
095878  
095879  
095880  
095881  
095882  
095883  
095884  
095885  
095886  
095887  
095888  
095889  
095890  
095891  
095892  
095893  
095894  
095895  
095896  
095897  
095898  
095899  
095900  
095901  
095902  
095903  
095904  
095905  
095906  
095907  
095908  
095909  
095910  
095911  
095912  
095913  
095914  
095915  
095916  
095917  
095918  
095919  
095920  
095921  
095922  
095923  
095924  
095925  
095926  
095927  
095928  
095929  
095930  
095931  
095932  
095933  
095934  
095935  
095936  
095937  
095938  
095939  
095940  
095941  
095942  
095943  
095944  
095945  
095946  
095947  
095948  
095949  
095950  
095951  
095952  
095953  
095954  
095955  
095956  
095957  
095958  
095959  
095960  
095961  
095962  
095963  
095964  
095965  
095966  
095967  
095968  
095969  
095970  
095971  
095972  
095973  
095974  
095975  
095976  
095977  
095978  
095979  
095980  
095981  
095982  
095983  
095984  
095985  
095986  
095987  
095988  
095989  
095990  
095991  
095992  
095993  
095994  
095995  
095996  
095997  
095998  
095999  
096000

The present invention provides in one embodiment a method of determining the genotype at a locus within genetic material obtained from a biological sample. In accordance with this method, the material is reacted at the locus to produce a first reaction value indicative of the presence of a given allele at the locus. There is formed a data set including the first reaction value. There is also established a set of one or more probability distributions; these distributions associate hypothetical reaction values with corresponding probabilities for each genotype of interest at the locus. The first reaction value is applied to each probability distribution to determine a measure of the conditional probability of each genotype of interest at the locus. The genotype is then determined based on these measures.

25

In accordance with a further embodiment of this method, the material at the locus is subject to a second reaction to produce a second reaction value independently indicative of the presence of a second allele at the locus. A second data set is formed and the second reaction value is included in the second data set. Each probability distribution associates a hypothetical pair of first and second reaction

30

5 values with a single probability of each genotype of  
interest. The first data set includes other reaction values  
obtained under conditions comparable to those under which  
the first reaction value was produced, and the second data  
set includes other reaction values obtained under conditions  
10 comparable to those under which the second reaction value  
was produced. Where, for example, there are two alleles of  
interest, the first reaction may be an assay for one allele  
and the second reaction may be a distinct assay for the  
other allele. The first and second data sets may include  
reaction values for the first and second reactions  
15 respectively, run under comparable conditions on other  
samples with respect to the same locus. Alternatively, or in  
addition, the data sets may include reaction values for  
reactions run under comparable conditions with respect to  
different loci within the same sample.  
20

In accordance with a further embodiment, the  
probability distributions may be determined iteratively. In  
this embodiment, each probability distribution is initially  
estimated. Each initial probability distribution is used to  
25 determine initial genotype probabilities using the reaction  
values in the data sets. The resulting data are then used to  
modify the initial probability distribution, so that the  
modified distribution more accurately reflects the reaction  
values in the data set. This procedure may be iterated a  
30 desired number of times to improve the probability  
distribution. In practice, we have generally found that a  
single iteration is sufficient.

05:41:37.07:000

5 The foregoing methods have been employed with success  
for automatic genotype determination based on assays using  
genetic bit analysis (GBA). In such a case, each allele may  
typically be a single specific nucleotide. In accordance  
with GBA, a reaction is designed to produce a value that is  
10 indicative of the presence of a specific allele at the locus  
within the genetic material. In GBA, the approach is  
typically to hybridize a specific oligonucleotide to the  
genetic material at the locus immediately adjacent to the  
nucleotide being interrogated. Next, DNA polymerase is  
15 applied in the presence of differentially labelled  
dideoxynucleoside triphosphates. The read-out steps detect  
the presence of one or more of the labels which have become  
covalently attached to the 3' end of the oligonucleotide.  
Details are provided in Theo R. Nikiforov et al. "Genetic  
20 Bit Analysis, a solid phase method for typing single  
nucleotide polymorphisms," 22 Nucleic Acids Research, No.  
20, 4167-4175 (1994), which is hereby incorporated herein by  
reference. However, the present invention is also applicable  
to other reaction systems for allele determination, such as  
25 allele-specific hybridization (ASH), sequencing by  
hybridization (CBH), oligonucleotide ligase assay (OLA), and  
allele-specific amplification, using either the ligase chain  
reaction (LCR) or the polymerase chain reactions (PCR). The  
alleles assayed may be defined, for example, by a single  
30 nucleotide, a pair of nucleotides, a restriction site, or  
(at least in part) by its length in nucleotides.

In another embodiment of the invention, there is

5 at the locus. Thereafter the distribution establishment arrangement modifies each initial probability distribution, so that each modified distribution more accurately reflects the reaction values stored in the storage means.

10 The term "reaction value" as used in this description and the following claims may refer either to a single numerical value or to a collection of numbers associated with a physical state produced by the reaction. In the GBA method described in the Nikiforov article referred to above, e.g., optical signals are produced that may be read as a single numerical value. Alternatively, e.g., an optical signal may be simplified over time, and the reaction value may be the collection of samples of such a signal. It is also possible to form a scanned image, of one or a series of optical signals generated by GBA or other reaction methods, and to digitize this image, so that a collection of pixel values in all or a portion of the image constitutes a reaction value.

#### Brief Description of the Drawings

25 The foregoing aspects of the invention will be more readily understood by reference to the following detailed description, taken with respect to the following drawings, in which:

30 Fig. 1 is a diagram of a device in accordance with a preferred embodiment of the invention;

Fig. 2 is a diagram of the logical flow in accordance

5 obtained under comparable conditions. A distribution  
establishment arrangement establishes a set of probability  
distributions, including at least one distribution,  
associating hypothetical reaction values with corresponding  
probabilities for each genotype of interest at the locus. A  
10 genotype calculation arrangement applies the first reaction  
value to each pertinent probability distribution to  
determine the conditional probability of each genotype of  
interest at the locus. A genotype determination arrangement  
determines the genotype based on data from the genotype  
5 calculation arrangement.

In a further embodiment, the device may determine the  
genotype at selected loci. In this embodiment, the reaction  
generation arrangement can produce a reaction value  
indicative of the presence of a given allele at each of the  
selected loci and the data set includes reaction values  
10 obtained with respect to each of the selected loci. The  
genotype calculation arrangement applies reaction values  
obtained with respect to each of the selected loci to each  
pertinent probability distribution.

25 In another further embodiment, the device may determine  
the genotype at a locus within genetic material from each of  
a plurality of samples. In this embodiment, the reaction  
generation arrangement can produce a reaction value  
indicative of the presence of a given allele at the locus of  
30 material obtained from each sample and the data set includes  
reaction values obtained with respect to each sample. The  
genotype calculation arrangement applies reaction values

5

10

25

arrangement then invokes the genotype calculation means to use each initial probability distribution to determine initial conditional probabilities for a genotype of interest.



5 provided a method of determining the genotype of a subject  
by reacting genetic material taken from the subject at  
selected loci. In this embodiment, each locus may be an  
identified single nucleotide or group of nucleotides, and  
there is produced with respect to each of the selected loci  
10 a reaction value indicative of the presence of a given  
allele at each of the selected loci. These reaction values  
are used to determine the genotype of the subject or  
alternatively a DNA sequence associated with a specific  
region of genetic material of the subject. (Indeed a set of  
15 genotypes for selected proximal loci may be used to specify  
a sequence of the genetic material.). In further embodiments,  
the loci are selected to provide one or more types of  
information concerning the subject, including inheritance of  
a trait, parentage, identity, and matching tissue with that  
20 of a donor. Alternatively, the loci may be spaced  
throughout the entire genome of subject to assist in  
characterizing the genome of the species of the subject.

25 In a further embodiment of the invention, there is  
provided a device for determining the genotype at a locus  
within genetic material obtained from a subject. The device  
of this embodiment has a reaction value generation  
arrangement for producing a first physical state,  
quantifiable as a first reaction value, indicative of the  
presence of a given allele at the locus, the value  
30 associated with reaction of the material at the locus. The  
device also has a storage arrangement for storing a data set  
including the first reaction value and other reaction values

5 with the embodiment of Fig. 1;

Fig. 3 is a graph of numeric reaction values (data) generated by the embodiment of Fig. 1 as well as the genotype determinations made by the embodiment from these data; and

10 Figs. 4-7 show probability distributions derived by the embodiment of Fig. 1 for three genotypes of interest (AA, AT, and TT) and a failure mode at a locus.

Fig. 8 is an example of the output of the device in Fig. 1.

#### Detailed Description of Specific Embodiments

The invention provides in preferred embodiments a method and device for genotype determination using genetic marker systems that produce allele-specific quantitative signals. An embodiment uses computer processing, employing computer software we developed and call "GetGenos", of data produced by a device we also developed to produce GBA data. The device achieves, among other things, the following:

- 25 • Fully automatic genotype determination from quantitative data. Off-line analysis of data pools is intended, although the software is fast enough to use interactively.
- Ability to examine many allele tests per DNA sample simultaneously. One genotype and confidence measure are produced from these data.
- 30 • A true probabilistic confidence measure (a LOD

5 score), properly calibrated, is produced for each genotype.

- Use of robust statistical methods: Noise reduction via selective data pooling and simultaneous search over points in a data pool, preventing bias.

- Maximal avoidance of arbitrary parameters, and thus  
10 insensitivity to great variation in input data. The small number of parameters that are required by the underlying statistical model are fit to the observed data, essentially using the data set as its own internal control.

- Flexibility for handling multiple data types.  
15 Essentially, only probability distribution calculations, described below, need to be calibrated to new data types. We expect that the invention may be applied to GBA, OLA, ASH, and RAPD-type markers.

Our current embodiment of the software is implemented  
20 in portable ANSI C, for easy integration into a custom laboratory information system. This code has been successfully run on:

- Macintosh
- Sun
- 25 • MS-DOS
- MS-Windows

In our current embodiment of the software, a number of consistency checks are performed for GBA data verification, using both the raw GBA values and the control wells.

30 Overall statistics for trend analysis and QC are computed. Brief "Genotype Reports" are generated, summarizing results for each data set, including failures. All data are output

5 in a convenient form for import into interactive statistical packages, such as DataDesk™. The current implementation is presently restricted to 2-allele tests in diploids - the situation with present GBA applications.

Referring to Fig. 1, there is shown a preferred  
10 embodiment of a device in accordance with the present invention. The device includes an optical detector 11 to produce reaction values resulting from one or more reactions. These reactions assay for one or more alleles in samples of genetic material. We have implemented the  
15 detector 11 using bichromatic microplate reader model 348 and microplate stacker model 83 from ICN Biomedical, Inc., P.O. Box 5023, Costa Mesa, California 92626. The microplates are in a 96 well format, and the reader accommodates 20  
20 microplates in a single processing batch. Accordingly the device of this embodiment permits large batch processing. The reactions in our implementation use GBA, as described above. The detector 11 is controlled by computer 12 to cause selected readout of reaction values from each well. The computer 12 is programmed to allow for multiple readout of  
25 the reaction value from a given well over a period of time. The values are stored temporarily in memory and then saved in database 14. Computer 13 accesses the database 14 over line 15 and processes the data in accordance with the procedure described below. Of course, computers 12 and 13  
30 and database 14 may be implemented by an integral controller and data storage arrangement. Such an arrangement could in fact be located in the housing of the optical detector 11.

5 In Fig. 2 is shown the procedure followed by computer  
13. The steps of this procedure are as follows:

Input Data: A set of data is loaded under step 21. In  
most applications, each experiment in the set should be  
testing (i) the same genetic marker, and (ii) the same set  
10 of alleles of that marker, using comparable biochemistry  
(e.g. the same reagent batches, etc.). Large data sets help  
smooth out noise, although the appropriate size of a data  
set depends on the allele frequencies (and thus the number  
of expected individuals of each genotypic class). Each data  
15 point in the input data may be thought of as an N-tuple of  
numeric values, where N is the number of signals collected  
from each DNA sample for this locus. (N will usually be the  
number of alleles tested at this marker, denoted A, except  
when repeated testing is used, in which case N may be  
20 greater than A).

Preprocess Data: Next the data are subject to  
preprocessing (step 22). An internal M-dimensional Euclidean  
representation of the input signals is produced, where each  
input datum (an N-tuple) is a point in M-space. Usually, M  
25 will be the same as N and the coordinates of the point will  
be the values of the input tuple, and thus the preprocessing  
will be trivial (although see the first paragraph of  
variations discussed). The Euclidean space may be  
non-linear, depending on the best available models of signal  
30 generation. (Completely mathematically equivalently, any  
non-linearity may be embodied in the initial probability  
distributions, described below.)

Fig. 3 illustrates preprocessed reaction values from step 22 for GBA locus 177-2 on 80 DNA samples. The X-axis indicates preprocessed reaction values for allele 1 (A) and the Y-axis indicates preprocessed reaction values for allele 2 (T). For clarity, the results of genotype determination are also indicated for each point: Triangles are TT genotype, diamonds are AA, circles are AT, and squares are failures (no signal).

Probability Distributions: Returning to Fig. 2, under step 22, initial probability distributions are established for the G possible genotypes. For example, in a random diploid population containing A tested alleles:

$$G = (A) + (A - 1) + \dots + 1 = \frac{A(A + 1)}{2} \quad (1)$$

The initial conditional probability for any hypothetical input datum (a point in M-space, denoted  $X_i$ ) and genotype (denoted  $g$ ) is defined as the prior probability of seeing the signal  $X_i$  assuming that  $g$  is the correct genotype of that datum. That is:

$$\begin{aligned} &\Pr(\text{signal } X_i \cdot \text{Genotype} = g), \\ &\text{where } X_i = (x_1^1 \dots x_1^M) \text{ and } g \in \{1 \dots G\} \end{aligned} \quad (2)$$

Figures 4 through 7 illustrate the initial probability distributions established for the data in figure 3. Probability distributions are indicated for the four

5 genotypic classes of interest, AA, AT, TT and No Signal, in  
Figs 4, 5, 6, and 7 respectively. The shading at each XY  
position indicates probability, with darker shades  
indicating increased probability for hypothetical data  
points with those X and Y reaction valves.

10 Exactly where these distributions come from is highly  
specific to the nature of the input data. The probability  
distributions can either be pre-computed at this step and  
stored as quantized data, or can be calculated on the fly as  
needed in step 23, below. The probability distributions may  
be fixed, or may be fit to the observed data or may be fit  
to assumed genotypes as determined by previous iterations of  
this algorithm. (See Additional Features below.)

Under step 23, we compute the conditional probability  
of each genotype. For each datum  $X_i$ , the above probabilities  
are collected into an overall conditional posterior  
probability of each genotype for that datum:

$$\begin{aligned} & \Pr(\text{Genotype} = g \mid \text{Signal } X_i) = \\ & \frac{\Pr(\text{Signal } X_i) \mid \text{Genotype} = g) \cdot \Pr(\text{Genotype} = g)}{\Pr(\text{Signal } X_i)} \quad (3) \end{aligned}$$

where

$\Pr(\text{Genotype} = g)$  is the prior probability of any datum  
having genotype  $g$ ;

$\Pr(\text{Signal } X_i)$  is the prior probability of the signal (a  
constant which may be ignored); and

$\Pr(\text{Signal } X_i) \cdot \text{Genotype} = g$  is the initial  
probability defined above.

Under step 24, we determine the select the genotype and compute the confidence score. For each datum, using the above posterior probabilities, we determine the most likely genotype assignment  $g'$  (the genotype with the highest posterior probability) and its confidence score. The confidence score  $C$  is simply the log of the odds ratio:

$$C = \log_{10} \frac{\Pr(\text{Genotype} = g' \mid \text{Signal } X_i)}{\sum_{\text{Genotypes } g} \Pr(\text{Genotype} = g \mid \text{Signal } X_i)} \quad (4)$$

It should be noted that this procedure is significant, among other reasons, because it permits determining a robust probabilistic confidence score associated with each genotype determination.

Under step 25, there may be employed adaptive fitting. A classic iterative adaptive fitting algorithm, such as Estimation-Maximization (E-M), may be used to increase the ability to deal with highly different input data sets and reduce noise sensitivity. In this case, the genotypes computed in step 24 are used to refit the distributions (from step 22). In step 25, a convergence test is performed, which may cause the program to loop back to step 23, but now using the new distributions.

As one example, an E-M search procedure may be used to maximize the total likelihood, that is, to find the maximally likely set of genotype assignments given the input data set. (The net likelihood may be calculated from the Bayesian probabilities, defined above.) For appropriate



5 likelihood calculations and probability distributions, the EM principle will guarantee that this algorithm always produces true-maximum-likelihood values, regardless of initial guess, and that it always converges.

10 Output Data: Under step 26, we output the results (genotypes and confidence scores) to the user or to a computer database. An example of such output is shown in Fig. 8.

#### Additional Features

15 Additional features may be incorporated into the above procedure. They may be integrated into the procedure either together or separately, and have all been implemented in a preferred embodiment.

20 Preprocessing: During steps 21 or 22, the data (either input tuples or spatial data points) may be preprocessed in order to reduce noise, using any one of many classical statistical or signal-processing techniques. Control data points may be used in this step. In fact, various types of signal filtering or normalizing may be applied at almost any step in the algorithm.

25 Fitting Probability Distributions: The probability distributions calculated in steps 22 and 23 may be fit to the input data - that is, each distribution may be a function of values which are in part calculated from the input data. For example, we may define the conditional probability of a signal point for some genotype to be a  
30 function of the distance between that point and the observed mean for that signal.

5        Using an Initial Genotype Guess: In step 22, either a  
 simple or heuristic algorithm may be used to produce an  
 initial genotype guess for each input data point. If a  
 fairly accurate guess can be produced, then the probability  
 distributions for each genotype may be fit to the subset of  
 10    the data assumed to be of that genotypic class. Another use  
 of a genotype guess is in initial input validity checks  
 and/or preprocessing (e.g. Step 22), before the remainder of  
 the algorithm is applied. To be useful, a guess need not  
 produce complete genotypic information, however.

15        Using a Null Genotypic Class: In steps 22 and all  
 further steps, one (or more) additional probability  
 distributions may be added to fit the data to the signals  
 one would expect to see if an experiment (e.g. that datum)  
 failed. E.g.,

$$\text{Pr}(\text{signal } X_i \cdot \text{Genotype} \cdot \{1 \dots G\})$$

20        The current implementation above is presently  
 restricted to  $M=2$  and  $N=2 \cdot R$ , where  $R$  is the number of  
 25    repeated tests of both alleles. We refer to the two alleles  
 as  $X$  and  $Y$ . The program understands the notion of "plates"  
 of data, a number of which make up a data set.

30        The Initial Guess Variation is employed to initially  
 fit distributions using the heuristic described below. The  
 Initial Guess is produced during the Preprocessing Step  
 which normalizes and background subtracts the input data,  
 and remove apparent outlier points as well. These steps are

5 performed separately for each allele's signal (i.e., 1  
dimensional analysis). In fact, this preprocessing is  
applied separately to each of the R repeated tests, and the  
test with the small total 2 dimension residual is chosen for  
10 use in further steps. Various other preprocessing and  
post-processing steps are employed for GBA data validation  
and QC. In particular, controls producing a known reaction  
value may be employed to assure integrity of the biochemical  
process. In a preferred embodiment, signals are assumed to  
be small positive numbers (between 0.0 and 5.0, with 0.0  
15 indicating that allele is likely not present in the sample,  
and larger values indicating that it may be.

To handle a wide range of input data signal strengths,  
the Adaptive Fitting Variation is employed. However, the  
program is hard-coded to perform exactly one or two  
20 interactions passes through step 25, which we find works  
well for existing GBA data.

The probability distributions we fit at present in  
steps 22 and 25 have as their only parameters (i) the ratio  
of the X and Y signals for heterozygotes, and (ii) the  
25 variance from the normalized means (0.0 negative for that  
allele, 1.0 for positive for that allele) along each axis  
separately. In fact, these later numbers are constrained to  
be at least a fixed minimum, which is rarely exceeded, so  
that the algorithm will work with very small quantities of  
30 data and will produce the behavior we want. These numbers  
are computed separately for each microtiter plate. The  
probability distributions are generated using the code

5 (written in C) attached hereto and incorporated herein by reference as Appendix A.

The Null-Class variant is used to provide genotypic class indicating No Signal.

10 Quality control may also be enhanced in a surprising manner using the procedures described here. In particular, the confidence score C of equation (4) serves as a robust indicator of the performance of the biochemical reaction system. For example, a downward trend in the confidence scores within a single batch or in successive batches may indicate deterioration of an important reagent or of a sample or miscalibration of the instrumentation.

15 Accordingly, in a preferred embodiment, the computer may be used to determine the presence of a downward trend in the confidence score over time calculated in reference to each of the following variables: the locus (is there a downward trend in the confidence score of a single locus relative to other loci tested?), the sample (is there a downward trend in the confidence score of a single sample relative to other samples tested?), plate (is there a downward trend in the confidence score of this plate relative to other plate?), and batch (relative to other batches). If a downward trend of statistical significance (using, for example a chi square test) is detected, an alarm condition is entered.

25 30 Because the confidence score is an accurate indication of the reliability of the reaction system and the genotype determination, a low confidence score associated with a

- 5 given determination is taken as indicating the need for retesting.

09618178 071800

## APPENDIX A

```
/* The probability distributions in Figures 4, 5, 6, and 7, respectively,
   correspond to the values of xx_prob, xy_prob, yy_prob, and ns_prob, for
   all possible values of the preprocessed reaction values (x_val and y_val)
   in the range of interest (0.0 to 3.0). */
```

```
/* We assume that the following global variables are set... */
double x_pos_mean, x_neg_mean, y_pos_mean, y_neg_mean;
double x_val, y_val;
```

```
/* And we set the following globals... */
double xx_prob, xy_prob, yy_prob, ns_prob;
```

```
#define POS_VARIANCE 0.25
#define POS_VARIANCE_INCREMENT 0.00
#define NEG_VARIANCE 0.05
#define NEG_VARIANCE_INCREMENT 0.10
#define HET_VARIANCE 0.10
#define HET_VARIANCE_INCREMENT 0.20
```

```
#define COND_NEG_PROB(val,given_val,val_mean) \
    normal_prob(val_mean-val,NEG_VARIANCE NEG_VARIANCE_INCREMENT*given_val)
```

```
#define COND_HET_PROB(val,given_val) \
    normal_prob(given_val-val,HET_VARIANCE + HET_VARIANCE_INCREMENT)
```

```
double normal_prob(deviation,sigma)
double deviation, sigma;
{
    double val=exp(-(deviation*deviation)/(2.0*sigma*sigma));
    return(val>=TINY_PROB ? val : TINY_PROB);
}
```

```
void compute_probs()
{
    double x_pos_prob, y_pos_prob, x_neg_prob, y_neg_prob;

    x_pos_prob=normal_prob((x_pos_mean-x_val), POS_VARIANCE);
    x_neg_prob=normal_prob((x_neg_mean-x_val), NEG_VARIANCE);
    y_pos_prob=normal_prob((y_pos_mean-y_val), POS_VARIANCE);
    y_neg_prob=normal_prob((y_neg_mean-y_val), NEG_VARIANCE);

    ns_prob=max(x_neg_prob * COND_NEG_PROB(y_val,x_val,y_neg_mean),
               y_neg_prob * COND_NEG_PROB(x_val,y_val,x_neg_mean));

    xx_prob=x_pos_prob * COND_NEG_PROB(y_val,x_val, y_neg_mean);

    yy_prob=y_pos_prob * COND_NEG_PROB(x_val,y_val, x_neg_mean);

    xy_prob= max(x_pos_prob * COND_HET_PROB(y_val,x_val),
               y_pos_prob * COND_HET_PROB(x_val,y_val));
}
```

00618178 071800